

**This Page Is Inserted by IFW Operations  
and is not a part of the Official Record**

## **BEST AVAILABLE IMAGES**

**Defective images within this document are accurate representations of  
the original documents submitted by the applicant.**

**Defects in the images may include (but are not limited to):**

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORLED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

673530

AK

P 00/008b 12/11/91  
Section 29 (1)  
Regulation 3.1 (2)

AUSTRALIA

Patents Act 1990

**NOTICE OF ENTITLEMENT**

(To be filed before acceptance)

We, **B A S F AKTIENGESELLSCHAFT** of, Carl-Bosch-Strasse 38, D-67056 Ludwigshafen, Germany being the applicant in respect of Application No. 45678/93 state the following:-

The Person nominated for the grant of the patent has entitlement from the actual inventors by virtue of their employment.

The person nominated for the grant of the patent is the applicant of the applications listed in the declaration under Article 8 of the PCT.

The basic applications listed on the request form are the first applications made in a Convention country in respect of the invention.

**B A S F AKTIENGESELLSCHAFT**

By our Patent Attorneys,

**WATERMARK PATENT & TRADEMARK ATTORNEYS**

27 June 1996

Louis C. Gebhardt

Registered Patent Attorney



(12) PATENT ABRIDGMENT (11) Document No AU-B-45678/93  
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No 673530

- (54) Title  
USE OF COMPOUNDS WHICH ABSORB AND/OR FLUORESC IN THE I/R RANGE AS MARKERS  
FOR LIQUIDS
- International Patent Classification(s)  
(51) G01N 021/64
- (21) Application No. : 45678/93 (22) Application Date 13.07.93
- (87) PCT Publication Number : WO94/02570
- (30) Priority Data
- | (31) Number | (32) Date | (33) Country |
|-------------|-----------|--------------|
| 4224301     | 23.07.92  | DE GERMANY   |
| 4243776     | 23.12.92  | DE GERMANY   |
| 4243774     | 23.12.92  | DE GERMANY   |
- (43) Publication Date : 14.02.94
- (44) Publication Date of Accepted Application : 14.11.96
- (71) Applicant(s)  
BASF AKTIENGESELLSCHAFT
- (72) Inventor(s)  
BERNHARD ALBERT; JUERGEN KIPPER; CHRISTOS VAMVAKARIS; KARIN HEIDRUN BECK;  
GERHARD WAGENBLAST
- (74) Attorney or Agent  
WATERMARK PATENT & TRADEMARK ATTORNEYS , Locked Bag 5, HAWTHORN VIC 3122
- (56) Prior Art Documents  
AU A1 55685/94 G01N 21/64  
US A 4735631  
US A 4209302
- (57) Claim

1. Use of a compound from the class of metal-free or metal-containing phthalocyanines, metal-free or metal-containing naphthalocyanines, nickel dithiolene complexes, aminium compounds of aromatic amines, methine dyes or azulenesquaric acid dyes which have their absorption maximum in the range from 600 to 1,200 nm and/or a fluorescence mixture in the range from 620 to 1,200 nm, as a marker for liquids.

4. A method for detecting a marker from the class of metal-free or metal-containing naphthalocyanines, metal-free or metal-containing naphthalocyanines, nickel dithiolene complexes, aminium compounds of aromatic amines, methine dyes or azulenesquaric acid dyes which have their fluorescence maximum in the range from 620 to 1,200 nm by exciting its fluorescence and so producing fluorescent light in the NIR spectral region.

5. A method as claimed in claim 4, wherein the fluorescence is excited by means of a semiconductor laser or a semiconductor diode.
6. A method as claimed in claim 4, wherein the fluorescence light is detected by means of a semiconductor detector.




(51) Internationale Patentklassifikation <sup>5</sup> : <b>C10L 1/10</b>		(11) Internationale Veröffentlichungsnummer: <b>WO 94/02570</b>
<b>A1</b>		(43) Internationales Veröffentlichungsdatum: <b>3. Februar 1994 (03.02.94)</b>
(21) Internationales Aktenzeichen: <b>PCT/EP93/01830</b> (22) Internationales Anmeldedatum: <b>13. Juli 1993 (13.07.93)</b> (30) Prioritätsdaten: P 42 24 301.7      23. Juli 1992 (23.07.92)      DE P 42 43 776.8      23. Dezember 1992 (23.12.92)      DE P 42 43 774.1      23. Dezember 1992 (23.12.92)      DE (71) Anmelder (für alle Bestimmungsstaaten ausser US): <b>BASF AKTIENGESELLSCHAFT [DE/DE]; Carl-Bosch-Strasse 38, D-67056 Ludwigshafen (DE).</b> (72) Erfinder; und (75) Erfinder/Anmelder (nur für US) : <b>ALBERT, Bernhard [DE/DE]; Riethburgstrasse 13, D-6701 Maxdorf (DE). KIPPER, Juergen [DE/DE]; Roonstrasse 26, D-7500 Karlsruhe 1 (DE). VAMVAKARIS, Christos [DE/DE]; Riedweg 6, D-6701 Kallstadt (DE). BECK, Karin, Heidrun [DE/DE]; Dackheimer Strasse 3, D-6700 Ludwigshafen (DE). WAGENBLAST, Gerhard [DE/DE]; Bachweg 8, D-6719 Weisenheim (DE).</b>		(81) Bestimmungsstaaten: <b>AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN, europäisches Patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b>  Veröffentlicht <i>Mit internationalem Recherchenbericht.</i>

673550

(54) Title: **USE OF COMPOUNDS WHICH ABSORB AND/OR FLUORESC IN THE I/R RANGE AS MARKERS FOR LIQUIDS**

(54) Bezeichnung: **VERWENDUNG VON IM IR-BEREICH ABSORBIERENDEN UND/ODER FLUORESZIERENDEN VERBINDUGEN ALS MARKIERUNGSMITTEL FÜR FLÜSSIGKEITEN**



(57) Abstract

The use of compounds from the class of the phthalocyanine, naphthalocyanine, nickel-dithiolene complexe, aminium compounds of aromatic amines, methine dyes or azulene quadratic acid dyes, the maximum absorption of which is in the 600 to 1,200 nm range and/or the maximum fluorescence of which is in the 620 to 1,200 nm range, as markers for liquids, a process for detecting markers in liquids and a suitable detector therefor.

(57) Zusammenfassung

Verwendung von Verbindungen aus der Klasse der Phthalocyanine, der Naphthalocyanine, der Nickel-Dithiolen-Komplexe, der Aminiumverbindungen von aromatischen Aminen, der Methinfarbstoffe oder der Azulenquadratsäurefarbstoffe, die ihr Absorptionsmaximum im Bereich von 600 bis 1 200 nm und/oder ein Fluoreszenzmaximum im Bereich von 620 bis 1 200 nm aufweisen, als Markierungsmittel für Flüssigkeiten, ein Verfahren zur Detektion von Markierungsmittel in Flüssigkeiten sowie ein dazu geeigneter Detektor.

Use of compounds which absorb and/or fluoresce  
in the IR region as markers for liquids

The present invention relates to the use of compounds from the class of metal-free or metal-containing phthalocyanines, metal-free or metal-containing naphthalocyanines, nickel dithiolene complexes, aminium compounds of aromatic amines, methine dyes or azulenesquaric acid dyes which have their absorption maximum in the range from 600 to 1,200 nm and/or a fluorescence maximum in the range from 620 to 1,200 nm, as markers (tracers) for liquids, to a method for detecting markers in liquids, and to a detector suitable for this purpose.

It is frequently necessary to mark liquids in order subsequently, for example when they are used, to re-detect the thus-marked liquids by means of suitable methods.

For example, heating oil can be differentiated from diesel oil in this way.


It is an object of the present invention to provide suitable compounds which are suitable as markers. The markers should have sufficiently strong absorption and/or fluorescence in the near infrared, so that detection of the absorption by means of conventional photometers which are sensitive in this range and/or of the fluorescence by means of conventional instruments after excitation with a suitable radiation source is possible.

We have found that this object is achieved by the compounds described in greater detail at the outset.

Metal-containing phthalocyanines or naphthalocyanines generally contain lithium (twice), magnesium, zinc, copper, nickel, VO, TiO or AlCl as the central atom.

Suitable phthalocyanines have, for example, the formula Ia






(Ia),

where

Me<sup>1</sup> is two hydrogen atoms, two lithium atoms or magnesium, zinc, copper, nickel, VO, TiO, AlCl or Si(OH),, at least 4 of the radicals R<sup>1</sup> to R<sup>16</sup>, independently of one another, have the formula W-X<sup>1</sup> in which W is a chemical bond, sulfur, imino, C<sub>1</sub>-C<sub>4</sub>-alkylimino or phenylimino, and X<sup>1</sup> is C<sub>1</sub>-C<sub>20</sub>-alkyl, which may be interrupted by from 1 to 4 oxygen atoms in ether function or substituted by phenyl, or are adamantyl or substituted or unsubstituted phenyl, and any remaining radicals R<sup>1</sup> to R<sup>16</sup> are hydrogen, halogen, hydroxysulfonyl or C<sub>1</sub>-C<sub>4</sub>-dialkylsulfamoyl.

Other suitable phthalocyanines have, for example, the formula Ib



(Ib),




where

$R^{17}$  and  $R^{18}$  or  $R^{18}$  and  $R^{19}$  or  $R^{19}$  and  $R^{20}$  together in each case have the formula  $X^2-C_6H_4-X^3$ , in which one of the two radicals  $X^2$  and  $X^3$  is oxygen and the other is imino or  $C_1-C_4$ -alkylimino, and

$R^{19}$  and  $R^{20}$  or  $R^{17}$  and  $R^{20}$  or  $R^{17}$  and  $R^{18}$ , independently of one another, are in each case hydrogen or halogen, and  $Me^1$  is as defined above.

Suitable naphthalocyanines have, for example, the formula II



(II),

where


$Y^1$ ,  $Y^2$ ,  $Y^3$ ,  $Y^4$ ,  $Y^5$ ,  $Y^6$ ,  $Y^7$  and  $Y^8$ , independently of one another, are each hydrogen, hydroxyl,  $C_1-C_{20}$ -alkyl or  $C_1-C_{20}$ -alkoxy, where the alkyl groups may each be interrupted by from 1 to 4 oxygen atoms in ether function and may be substituted by phenyl,

$Y^9$ ,  $Y^{10}$ ,  $Y^{11}$  and  $Y^{12}$ , independently of one another, are each hydrogen,  $C_1-C_{20}$ -alkyl or  $C_1-C_{20}$ -alkoxy, where the alkyl groups may each be interrupted by from 1 to 4 oxygen atoms in ether function, or are halogen, hydroxysulfonyl or  $C_1-C_4$ -dialkylsulfamoyl, and

$Me^2$  is two hydrogen atoms, two lithium atoms, magnesium, zinc, copper, nickel, VO, TiO, AlCl or








where


$\text{Y}^{17}$  and  $\text{Y}^{18}$ , independently of one another, are each hydroxyl,  $\text{C}_1\text{-C}_{20}$ -alkoxy,  $\text{C}_1\text{-C}_{20}$ -alkyl,  $\text{C}_2\text{-C}_{20}$ -alkenyl,  $\text{C}_3\text{-C}_{20}$ -alkenyloxy or



5 in which  $\text{Y}^{19}$  is  $\text{C}_1\text{-C}_{20}$ -alkyl,  $\text{C}_2\text{-C}_{20}$ -alkenyl or  $\text{C}_4\text{-C}_{20}$ -alkadienyl and  $\text{Y}^{20}$  and  $\text{Y}^{21}$ , independently of one another, are each  $\text{C}_1\text{-C}_{12}$ -alkyl,  $\text{C}_2\text{-C}_{12}$ -alkenyl or as defined for  $\text{OY}^{19}$  above.


10 Of particular interest here are naphthalocyanines of the formula II in which at least one of  $\text{Y}^1$  to  $\text{Y}^8$  is not hydrogen.

Suitable nickel dithiolene complexes have, for example, the formula III




where

15  $\text{L}^1$ ,  $\text{L}^2$ ,  $\text{L}^3$  and  $\text{L}^4$ , independently of one another, are each  $\text{C}_1\text{-C}_{20}$ -alkyl, which may be interrupted by from 1 to 4 oxygen atoms in ether function, phenyl,  $\text{C}_1\text{-C}_{20}$ -alkylphenyl,  $\text{C}_1\text{-C}_{20}$ -alkoxyphenyl, where the alkyl groups may  
20 function, or  $\text{L}^1$  and  $\text{L}^2$  and/or  $\text{L}^3$  and  $\text{L}^4$  in each case together are




Suitable aminium compounds have, for example, the formula IV



where


5  $Z^1$ ,  $Z^2$ ,  $Z^3$  and  $Z^4$ , independently of one another, are each  $C_1$ - $C_{20}$ -alkyl, which may be interrupted by from 1 to 4 oxygen atoms in ether function,  $C_1$ - $C_{20}$ -alkanoyl or



10 where  $Z^6$  is hydrogen,  $C_1$ - $C_{20}$ -alkyl, which may be interrupted by from 1 to 4 oxygen atoms in ether function, or  $C_1$ - $C_{20}$ -alkanoyl,  $Z^7$  is hydrogen or  $C_1$ - $C_{20}$ -alkyl, which may be interrupted by from 1 to 4 oxygen atoms in ether function, and  $Z^8$  is hydrogen,  $C_1$ - $C_{20}$ -alkyl, which may be interrupted by from 1 to 4 oxygen atoms in ether function, or halogen, and  $An^\ominus$  is the equivalent of an anion.

15 Suitable methine dyes have, for example, the formula V






in which the rings A and B, independently of one another, may each be benzo-fused and/or substituted,

E<sup>1</sup> and E<sup>2</sup>, independently of one another, are each oxygen, sulfur, imino, -C(CH<sub>3</sub>)<sub>2</sub>- or -CH=CH-,

5 D is



where E<sup>3</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, chlorine or bromine and E<sup>4</sup> is hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl,


10 Q<sup>1</sup> and Q<sup>2</sup>, independently of one another, are each phenyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>1</sub>-C<sub>12</sub>-alkyl, which may be interrupted by from 1 to 3 oxygen atoms in ether function and may be substituted by hydroxyl, chlorine, bromine, carboxyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, acryloyloxy, methacryloyloxy, hydroxysulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkanoylamino, C<sub>1</sub>-C<sub>6</sub>-alkylcarbamoyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbamoyloxy or G<sup>\*</sup>(K)<sub>n</sub>, where G is nitrogen or  
15 phosphorus and K is phenyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl or C<sub>1</sub>-C<sub>12</sub>-alkyl,

An<sup>o</sup> is the equivalent of an anion, and

n is 1, 2 or 3.

20 Suitable azulenescuaric acid dyes have, for example, the formula VI





(VI).

where

J is C<sub>1</sub>-C<sub>12</sub>-alkylene,

T<sup>1</sup> is hydrogen, halogen, amino, hydroxyl, C<sub>1</sub>-C<sub>12</sub>-alkoxy, phenyl, substituted phenyl, carboxyl, C<sub>1</sub>-C<sub>12</sub>-alkoxy-carbonyl, cyano, -NT<sup>6</sup>-CO-T<sup>7</sup>, -CO-NT<sup>6</sup>T<sup>7</sup> or -O-CO-NT<sup>6</sup>T<sup>7</sup>, where T<sup>6</sup> and T<sup>7</sup>, independently of one another, are each hydrogen, C<sub>1</sub>-C<sub>12</sub>-alkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, phenyl, 2,2,6,6-tetramethylpiperidin-4-yl or cyclohexylamino-carbonyl, and

T<sup>2</sup>, T<sup>3</sup>, T<sup>4</sup> and T<sup>5</sup>, independently of one another, are each hydrogen or C<sub>1</sub>-C<sub>12</sub>-alkyl, which may be substituted by halogen, amino, C<sub>1</sub>-C<sub>12</sub>-alkoxy, phenyl, substituted phenyl, carboxyl, C<sub>1</sub>-C<sub>12</sub>-alkoxycarbonyl or cyano, with the proviso that, if T<sup>5</sup> is hydrogen, the ring positions of the substituents J-T<sup>1</sup> and T<sup>4</sup> within an azulene ring may also be exchanged for one another on one or both azulene rings.

All the alkyl, alkylene and alkenyl radicals occurring in the above formulae may be either straight-chain or branched.

In the formulae Ia, II, III and IV, suitable C<sub>1</sub>-C<sub>20</sub>-alkyl radicals, which may be interrupted by from 1 to 4 oxygen atoms in ether function, are, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, tert-pentyl, hexyl, 2-methylpentyl, heptyl, octyl, 2-ethylhexyl, isooctyl, nonyl, isononyl, decyl, isodecyl, undecyl, dodecyl, tridecyl, 3,5,5,7-tetramethylnonyl,



isotridecyl (the above terms isooctyl, isononyl, isodecyl and isotridecyl are trivial names and originate from the alcohols obtained from the oxo synthesis - cf. in this respect Ullmanns Encyklopädie der technischen Chemie, 4th Edition, Volume 7, pages 215-217, and Volume 11, pages 435 and 436), tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, 2-methoxyethyl, 2-ethoxyethyl, 2-propoxyethyl, 2-isopropoxyethyl, 2-butoxyethyl, 2- and 3-methoxypropyl, 2- and 3-ethoxypropyl, 2- and 3-propoxypropyl, 2- and 3-butoxypropyl, 2- and 4-methoxybutyl, 2- and 4-ethoxybutyl, 2- and 4-propoxybutyl, 2- and 4-butoxybutyl, 3,6-dioxaheptyl, 3,6-dioxaoctyl, 4,8-dioxanonyl, 3,7-dioxaoctyl, 3,7-dioxanonyl, 4,7-dioxaoctyl, 4,7-dioxanonyl, 4,8-dioxadecyl, 3,6,8-trioxadecyl, 3,6,9-trioxaundecyl, 3,6,9,12-tetraoxatridecyl and 3,6,9,12-tetraoxatetradecyl.

In the formulae I and II, suitable phenyl-substituted  $C_1$ - $C_{20}$ -alkyl radicals are, for example, benzyl and 1- and 2-phenylethyl.

In the formulae II, III and IV, suitable  $C_1$ - $C_{20}$ -alkoxy radicals, which may be interrupted by from 1 to 4 oxygen atoms in ether function, are, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, pentoxy, hexyloxy, heptyloxy, octyloxy, 2-ethylhexyloxy, isooctyloxy, nonyloxy, isononyloxy, decyloxy, isodecyloxy, undecyloxy, dodecyloxy, tridecyloxy, isotridecyloxy, tetradecyloxy, pentadecyloxy, hexadecyloxy, heptadecyloxy, octadecyloxy, nonadecyloxy, eicosyloxy, 2-methoxyethoxy, 2-ethoxyethoxy, 2-propoxyethoxy, 2-isopropoxyethoxy, 2-butoxyethoxy, 2- and 3-methoxypropoxy, 2- and 3-ethoxypropoxy, 2- and 3-propoxypropoxy, 2- and 3-butoxypropoxy, 2- and 4-methoxybutoxy, 2- and 4-ethoxybutoxy, 2- and 4-propoxybutoxy, 2- and 4-butoxybutoxy, 3,6-dioxaheptyloxy, 3,6-dioxaoctyloxy, 4,8-dioxanonyloxy, 3,7-dioxaoctyloxy, 3,7-dioxanonyloxy, 4,7-dioxaoctyloxy, 4,7-dioxanonyloxy, 4,8-dioxadecyloxy, 3,6,8-trioxadecyloxy, 3,6,9-trioxaundecyloxy,



3,6,9,12-tetraoxatridecyloxy and 3,6,9,12-tetraoxatetradecyloxy.

5 In the formula II, suitable phenyl-substituted  $C_1$ - $C_{20}$ -alkoxy radicals are, for example, benzyloxy and 1- and 2-phenylethoxy.

10 In the formulae Ia, III and VI, suitable substituted phenyl radicals are, for example, phenyl which is substituted by  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkoxy, hydroxyl or halogen. In general, there may be from 1 to 3 substituents.

Halogen in the formulae Ib, II, IV and VI is, for example, fluorine, chlorine or bromine.

15 The radicals W in the formula Ia and  $X^2$  and  $X^3$  in the formula Ib are, for example, methylimino, ethylimino, propylimino, isopropylimino or butylimino.

The radicals  $R^1$  to  $R^{16}$  in the formula Ia and  $Y^9$  to  $Y^{12}$  in the formula II are, for example, dimethylsulfamoyl, diethylsulfamoyl, dipropylsulfamoyl, dibutylsulfamoyl or N-methyl-N-ethylsulfamoyl.

20  $C_2$ - $C_{20}$ -alkenyl and  $C_4$ - $C_{20}$ -alkandienyl in the formula II are, for example, vinyl, allyl, prop-1-en-1-yl, methallyl, ethallyl, but-3-en-1-yl, pentenyl, penta-  
dienyl, hexadienyl, 3,7-dimethylocta-1,6-dien-1-yl,  
undec-10-en-1-yl, 6,10-dimethylundeca-5,9-dien-2-yl,  
25 octadec-9-en-1-yl, octadeca-9,12-dien-1-yl,  
3,7,11,15-tetramethylhexadec-1-en-3-yl or  
eicos-9-en-1-yl.

30  $C_3$ - $C_{20}$ -alkenyloxy in the formula II is, for example, allyloxy, methallyloxy, but-3-en-1-yloxy, undec-10-en-1-yloxy, octadec-9-en-1-yloxy or eicos-9-en-1-yloxy.

$Z^6$  in the formula IV is, for example, formyl, acetyl, propionyl, butyryl, isobutyryl, pentanoyl, hexanoyl, heptanoyl, octanoyl or 2-ethylhexanoyl.

35 If the rings A and/or B in the formula V are substituted, suitable substituents are, for example,  $C_1$ - $C_6$ -alkyl, phenyl- $C_1$ - $C_6$ -alkoxy, phenoxy, halogen,



hydroxyl, amino, C<sub>1</sub>-C<sub>6</sub>-mono- or dialkylamino or cyano. The rings are generally monosubstituted to trisubstituted.

The radicals E<sup>1</sup>, E<sup>4</sup>, Q<sup>1</sup> and Q<sup>2</sup> in the formula V are, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, pentyl, isopentyl, neopentyl, tert-pentyl or hexyl.

The radicals Q<sup>1</sup> and Q<sup>2</sup> are furthermore, for example, hexyl, 2-methylpentyl, heptyl, octyl, 2-ethylhexyl, isooctyl, nonyl, isononyl, decyl, isodecyl, undecyl, dodecyl, cyclopentyl, cyclohexyl, 2-methoxyethyl, 2-ethoxyethyl, 2- or 3-methoxypropyl, 2- or 3-ethoxypropyl, 2-hydroxyethyl, 2- or 3-hydroxypropyl, 2-chloroethyl, 2-bromoethyl, 2- or 3-chloropropyl, 2- or 3-bromopropyl, 2-carboxyethyl, 2- or 3-carboxypropyl, 2-methoxycarbonyl-ethyl, 2-ethoxycarbonyl-ethyl, 2- or 3-methoxycarbonylpropyl, 2- or 3-ethoxycarbonylpropyl, 2-acryloyloxyethyl, 2- or 3-acryloyloxypropyl, 2-methacryloyloxyethyl, 2- or 3-methacryloyloxypropyl, 2-hydroxysulfonyl-ethyl, 2- or 3-hydroxysulfonylpropyl, 2-acetylamin-ethyl, 2- or 3-acetylaminopropyl, 2-methylcarbamoyl-ethyl, 2-ethylcarbamoyl-ethyl, 2- or 3-methylcarbamoylpropyl, 2- or 3-ethylcarbamoylpropyl, 2-methylcarbamoyloxyethyl, 2-ethylcarbamoyloxyethyl, 2- or 3-methylcarbamoyloxypropyl, 2- or 3-ethylcarbamoyloxypropyl, 2-(trimethylammonium)ethyl, 2-(triethylammonium)ethyl, 2- or 3-(trimethylammonium)propyl, 2- or 3-(triethylammonium)propyl, 2-(triphenylphosphonium)ethyl or 2- or 3-(triphenylphosphonium)propyl.

An<sup>o</sup> in the formulae IV and V is derived, for example, from anions of organic or inorganic acids. Particular preference is given to, for example, methane-sulfonate, 4-methylbenzenesulfonate, acetate, tri-fluoroacetate, heptafluorobutyrate, chloride, bromide, iodide, perchlorate, tetrafluoroborate, nitrate, hexafluorophosphate and tetraphenylborate.

The radicals J in the formula VI are, for example, methylene, ethylene, 1,2- or 1,3-propylene,

1,2-, 1,3-, 2,3- or 1,4-butylene, pentamethylene, hexamethylene, heptamethylene, octamethylene, nonamethylene, decamethylene, undecamethylene or dodecamethylene.

The radicals T<sup>2</sup>, T<sup>3</sup>, T<sup>4</sup> and T<sup>5</sup> in the formula VI are, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, tert-pentyl, 2-methylbutyl, hexyl, 2-methylpentyl, heptyl, octyl, 2-ethylhexyl, isooctyl, nonyl, isononyl, decyl, undecyl, dodecyl, fluoromethyl, chloromethyl, difluoromethyl, trifluoromethyl, trichloromethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromomethyl, 1,1,1-trifluoroethyl, heptafluoropropyl, 4-chlorobutyl, 5-fluoropentyl, 6-chlorohexyl, cyanomethyl, 2-cyanoethyl, 3-cyanopropyl, 2-cyanobutyl, 4-cyanobutyl, 5-cyanopentyl, 6-cyanoethyl, 2-aminoethyl, 2-aminopropyl, 3-aminopropyl, 2-aminobutyl, 4-aminobutyl, 5-aminopentyl, 6-aminohexyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 2-hydroxybutyl, 4-hydroxybutyl, 5-hydroxypentyl, 6-hydroxyhexyl, 2-methoxyethyl, 2-ethoxyethyl, 2-propoxyethyl, 2-isopropoxyethyl, 2-butoxyethyl, 2-methoxypropyl, 2-ethoxypropyl, 3-ethoxypropyl, 4-ethoxybutyl, 4-isopropoxybutyl, 5-ethoxypentyl, 6-methoxyhexyl, benzyl, 1-phenylethyl, 2-phenylethyl, 4-chlorobenzyl, 4-methoxybenzyl, 2-(4-methylphenyl)ethyl, carboxymethyl, 2-carboxyethyl, 3-carboxypropyl, 4-carboxybutyl, 5-carboxypentyl, 6-carboxyhexyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, 2-methoxycarbonylethyl, 2-ethoxycarbonylethyl, 3-methoxycarbonylpropyl, 3-ethoxycarbonylpropyl, 4-methoxycarbonylbutyl, 4-ethoxycarbonylbutyl, 5-methoxycarbonylpentyl, 5-ethoxycarbonylpentyl, 6-methoxycarbonylhexyl or 6-ethoxycarbonylhexyl.

T<sup>1</sup> in the formula VI is, for example, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl, pentoxycarbonyl, isopentoxycarbonyl, neopentoxycarbonyl, tert-pentoxycarbonyl, hexyloxycarbonyl, heptyloxycarbonyl, octyloxy-






carbonyl, isooctyloxycarbonyl, nonyloxycarbonyl, isononyloxycarbonyl, decyloxycarbonyl, isodecyloxycarbonyl, undecyloxycarbonyl, dodecyloxycarbonyl, methoxy, thoxy, propoxy, isopropoxy, butoxy, isobutoxy, pentoxy, hexyloxy, acetyl amino, carbamoyl, mono- or dimethylcarbamoyl, mono- or diethylcarbamoyl, monocyclohexylcarbonyl, phenylcarbamoyl, dimethylcarbamoyloxy or diethylcarbamoyloxy.


Preference is given to the use according to the invention of compounds originating from the class of metal-free or metal-containing naphthalocyanines.

Emphasis should be placed on the use according to the invention of naphthalocyanines of the formula IIa



where

$Y^1, Y^2, Y^3, Y^4, Y^5, Y^6, Y^7$  and  $Y^8$ , independently of one another, are each hydrogen, hydroxyl,  $C_1-C_4$ -alkyl or  $C_1-C_{20}$ -alkoxy, and  $Me^2$  is two hydrogen atoms, two lithium atoms, magnesium, zinc, copper, nickel, VO, AlCl or



where  $R^{19}$  is  $C_1-C_{11}$ -alkyl or  $C_{10}-C_{20}$ -alkadienyl and  $Y^{20}$  and  $Y^{21}$ , independently of one another, are each  $C_1-C_{11}$ -alkyl or




C<sub>2</sub>-C<sub>4</sub>-alkenyl.

Particular emphasis should be placed on the use according to the invention of naphthalocyanines of the formula IIa, where Y<sup>1</sup>, Y<sup>2</sup>, Y<sup>3</sup>, Y<sup>4</sup>, Y<sup>5</sup>, Y<sup>6</sup>, Y<sup>7</sup> and Y<sup>8</sup>, independently of one another, are each hydroxyl or C<sub>1</sub>-C<sub>20</sub>-alkoxy, in particular C<sub>1</sub>-C<sub>10</sub>-alkoxy. The alkoxy radicals here may be identical or different.

Particular emphasis should furthermore be placed on the use according to the invention of naphthalocyanines of the formula IIa where Me<sup>2</sup> is two hydrogen atoms.

Emphasis should furthermore be placed on the use according to the invention of nickel dithiolene complexes of the formula III, where L<sup>1</sup>, L<sup>2</sup>, L<sup>3</sup> and L<sup>4</sup>, independently of one another, are each phenyl, C<sub>1</sub>-C<sub>20</sub>-alkylphenyl, C<sub>1</sub>-C<sub>20</sub>-alkoxyphenyl or hydroxyl- and C<sub>1</sub>-C<sub>20</sub>-alkyl substituted phenyl, or L<sup>1</sup> and L<sup>2</sup> and L<sup>3</sup> and L<sup>4</sup> are in each case together



Particular emphasis should furthermore be placed on the use according to the invention of nickel dithiolene complexes of the formula III where L<sup>1</sup> and L<sup>4</sup> are each phenyl and L<sup>2</sup> and L<sup>3</sup> are each 4-[C<sub>2</sub>H<sub>5</sub>-C(CH<sub>3</sub>)<sub>2</sub>]-C<sub>6</sub>H<sub>4</sub>.

The phthalocyanines of the formula Ia are known per se and are described, for example, in DE-B-1 073 739 or EP-A-155 780 or can be obtained by methods known per se, as used in the preparation of phthalocyanines or naphthalocyanines and as described, for example, in F.H. Moser, A.L. Thomas, "The Phthalocyanines", CRC Press, Boca Rota, Florida, 1983, or J. Am. Chem. Soc. 106 (1984), 7404 to 7410. The phthalocyanines of the formula



Ib are likewise known per se and are described, for example, in EP-A-155 780 or can be obtained by the above prior-art methods (Moser, J. Am. Chem. Soc.).

5       The naphthalocyanines of the formula II are likewise known per se and are described, for example, in EP-A-336 213,       EP-A-358 080,       GB-A-2,168,372 or GB-A-2,200,650, or can be obtained by the above prior-art methods (Moser, J. Am. Chem. Soc.).

10       The nickel dithiolene complexes of the formula III are likewise known per se and are described, for example, in EP-A-192 215.

15       The aminium compounds of the formula IV are likewise known per se and are described, for example, in US-A-3,484,467 or can be obtained by the methods mentioned therein.

      The methine dyes of the formula V are likewise known per se and are described, for example, in EP-A-464 543 or can be obtained by the methods described therein.

20       The azulenesquaric acid dyes of the formula VI are likewise known per se and are described, for example, in EP-A-310 080 or US-A-4,990,649 or can be obtained by the methods described therein.

25       Suitable solvents which can be marked according to the invention by means of the compounds described in greater detail above are, in particular, organic liquids, for example alcohols, such as methanol, ethanol, propanol, isopropanol, butanol, isobutanol, sec-butanol, pentanol, isopentanol, neopentanol and hexanol, glycols, 30       such as 1,2-ethylene glycol, 1,2- and 1,3-propylene glycol, 1,2-, 2,3- and 1,4-butylene glycol, di- and triethylene glycol and di- and tripropylene glycol, ethers, such as methyl tert-butyl ether, 1,2-ethylene glycol mono- and dimethyl ether, 1,2-ethylene glycol 35       mono- and diethyl ether, 3-methoxypropanol, 3-isopropoxypropanol, tetrahydrofuran and dioxane, ketones, such as acetone, methyl ethyl ketone and diacetone alcohol,



esters, such as methyl acetate, ethyl acetate, propyl acetate and butyl acetate, aliphatic and aromatic hydrocarbons, such as pentane, hexane, heptane, octane, isooctane, petroleum ether, toluene, xylene, ethylbenzene, tetralin, decalin, dimethylnaphthalene, white spirit, mineral oil, such as gasoline, kerosene, diesel oil and heating oil, natural oils, such as olive oil, soya oil and sunflower oil, and natural or synthetic engine oils, hydraulic oils and transmission oils, for example vehicle engine oil and sewing machine oil, and brake fluids.

The abovementioned compounds are particularly advantageously used for marking mineral oils in which labeling is simultaneously required, for example for tax reasons. In order to keep costs of the labeling low, the usual aim is to use very dilutable dyes for the coloring. However, even dyes of high tinctorial strength cannot be detected purely visually in high dilution in mineral oils.

For this reason, it is of particular advantage to use markers which have their absorption maximum in the range from 600 to 1,200 nm and/or fluoresce in the range from 620 to 1,200 nm, since they can easily be detected using suitable instruments.

For labeling liquids, but in particular mineral oil, the abovementioned compounds are generally used in the form of solutions. Suitable solvents are preferably aromatic hydrocarbons, such as toluene and xylene. In order to avoid any excessively high viscosity of the resultant solutions, a concentration of the compound which absorbs and/or fluoresces in the IR of from 2 to 50 % by weight, based on the solution, is generally selected.

The present invention furthermore provides a process for detecting markers in liquids, which comprises detecting the marker from its fluorescence in the NIR (near infrared) spectral region.



The fluorescence of the markers present in the liquids is advantageously excited by means of a semiconductor laser or a semiconductor diode. It is particularly favorable to use a semiconductor laser or a semiconductor diode having a wavelength of maximum emission in the spectral range from  $\lambda_{\text{max}}-100$  nm to  $\lambda_{\text{max}}+20$  nm, where  $\lambda_{\text{max}}$  is the wavelength of the absorption maximum of the marker. The wavelength of maximum emission is in the range from 620 to 1,200 nm.

The fluorescence light produced in this way is advantageously detected by means of a semiconductor detector, in particular a silicon photodiode or a germanium photodiode.

Detection is particularly advantageous if the detector is preceded by an interference filter and/or a cut-off filter (with a short-wave transmission cut-off limit in the range from  $\lambda_{\text{max}}$  to  $\lambda_{\text{max}}+80$  nm) and/or a polarizer.

The abovementioned compounds make it very simple to detect marked liquids, even if the marker substances are only present in a concentration of approximately 0.1 ppm (detection by absorption) or approximately 5 ppb (detection by fluorescence).

The present invention furthermore provides an instrument for carrying out the process according to the invention (= detector), the detector comprising an NIR light source (semiconductor laser or semiconductor diode), one or more optical filters, an NIR polarizer and a photodetector (silicon photodiode or germanium photodiode) and, if desired, optical fibers or optical fiber bundles.

The examples below are intended to illustrate the invention in greater detail.


General procedure for detecting the markers.

#### I. Detection by absorption in the IR region



EXAMPLE 1

Sufficient dy of the formula



was dissolved in diesel fuel to give a solution containing 1,000 ppm of dye.

5 This solution was diluted further in steps, and its absorption in the NIR region measured against a pure diesel fuel using a commercially available spectrometer (1 cm cell).

10	Dye content in diesel fuel [ppm]	Absorption	Absorption maximum [nm]
	100	>> 3	-
	50	3.05	844.0
	20	2.81	854.0
	10	2.10	860.4
15	1	0.27	860.0

Similarly favorable results are achieved when naphthalocyanines of the above formula (where R = n-C<sub>8</sub>H<sub>17</sub>, or n-C<sub>12</sub>H<sub>25</sub>) or the dyes listed below are used for the marking.

20 Dye 2  
Hexadecaphenylthiocopper phthalocyanine


Dye 3  
Tetradecaphenylthiocopper phtahlocyanine



Dye 4

Tetradecadodecylthiocopper phthalocyanine

Dye 5




Dye 6

5 Hexadeca(4-tert-butylphenylthio)copper phthalocyanine


Dye 7

NcSi[-O-Si(CH<sub>3</sub>)<sub>2</sub>-O-C<sub>12</sub>H<sub>25</sub>]<sub>2</sub>


Dye 8




Dye 9











Dye 10




Dyes 11 to 15





Farb- stoff- Nr.	L <sup>1</sup>	L <sup>2</sup>	L <sup>3</sup>	L <sup>4</sup>
11		C <sub>6</sub> H <sub>5</sub>		C <sub>6</sub> H <sub>5</sub>
12		C <sub>6</sub> H <sub>5</sub>		C <sub>6</sub> H <sub>5</sub>
13	C <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub>
14		C <sub>6</sub> H <sub>5</sub>		C <sub>6</sub> H <sub>5</sub>
15				


Dyes 15 to 18



Farbstoff Nr.	Z	An <sup>⊖</sup>
16	C <sub>4</sub> H <sub>9</sub>	NO <sub>3</sub> <sup>⊖</sup>
17	C <sub>2</sub> H <sub>5</sub>	NO <sub>3</sub> <sup>⊖</sup>
18	C <sub>4</sub> H <sub>9</sub>	BF <sub>4</sub> <sup>⊖</sup>

Dyes 19 to 23


5



Farbstoff Nr.	Q <sup>1</sup>	Q <sup>2</sup>	Q <sup>3</sup>	Q <sup>4</sup>	An <sup>⊖</sup>
19	CH <sub>3</sub>	CH <sub>3</sub>	Cl	Cl	I <sup>⊖</sup>
20	CH <sub>3</sub>	CH <sub>3</sub>	H	H	I <sup>⊖</sup>
21	$\begin{array}{c} \text{O} \\    \\ \text{C}_2\text{H}_4\text{OCNHC}(\text{CH}_3)_3 \end{array}$	$\begin{array}{c} \text{O} \\    \\ \text{C}_2\text{H}_4\text{OCNHC}(\text{CH}_3)_3 \end{array}$	H	H	I <sup>⊖</sup>
22	$\begin{array}{c} \text{O} \\    \\ \text{C}_2\text{H}_4\text{CNHC}_6\text{H}_{13} \end{array}$	$\begin{array}{c} \text{O} \\    \\ \text{C}_2\text{H}_4\text{CNHC}_6\text{H}_{13} \end{array}$	H	H	ClO <sub>4</sub> <sup>⊖</sup>
23	C <sub>3</sub> H <sub>6</sub> SO <sub>3</sub> <sup>⊖</sup>	C <sub>3</sub> H <sub>6</sub> SO <sub>3</sub> H	H	H	Betain



## Dye 24



## II. Detection by fluorescence in the NIR region

Figure 1 shows the diagrammatic structure of the detector.

5        The marker fluorescence is excited using the emission from a commercial semiconductor diode laser. The parallel laser beam is shone onto the sample in a 1 cm cell. In order to double the excitation intensity, the transmitted light beam is reflected by a mirror and again  
10        shone through the sample.

15        The fluorescence light is imaged by means of optical elements (lens system) onto the detector, a silicon photodiode. The light emitted at the rear is likewise thrown onto the silicon photodiode by a concave mirror.


20        In order to separate the unwanted light (scattered excitation light) from the fluorescence light, cut-off and/or interference filters and/or a polarizer (NIR polarization foil) are used.

      The optimization of the polarizer is selected so that the direction of maximum transmission is perpendicular to the polarization plane of the excitation light.



Example 25

Sufficient dye of the formula



was dissolved in diesel fuel to give a stock solution containing 219 ppb of dye. Further solutions were prepared therefrom by dilution with diesel fuel.

These solutions were measured in accordance with general procedure II using the following equipment settings.

Excitation: GaAlAs semiconductor diode laser having a wavelength of 813 nm; CW output 7 mW.

Filter: Long-pass interference filter 850 nm (Corion).

Photodetector: Silicon photodiode having an area of 1 cm<sup>2</sup> (UDT). The photocurrent was measured using a current/voltage converter (UDT, model 350).


The results are shown in the table below.

Dye content in the diesel fuel [ppb]	Absorbance at $\lambda_{max}$	Fluorescence signal (in scale units)
219	0.05	2366
43.7	0.01	451
8.75	0.002	106
1.75	0.0004	40
0	0.0	20

This gives a lower detection limit for the marker by fluorescence of about 5 ppb.



Similarly favorable results are achieved when naphthalocyanines of the above formula (where  $R = n-C_5H_{11}$  or  $n-C_{12}H_{25}$ ) or the dyes listed below are used for the marking.



( $R = n-C_5H_{11}$  or  $n-C_{12}H_{25}$ )


5 Dye 26  
Me = 2H

Dye 27  
Me = Zn


10 Dye 28  
Me = AlCl

Dye 29  
 $NcSi[-O-Si(CH_3)_2-O-C_{12}H_{25}]_2$


Dye 30




Dye 31




Dye 32




Dye 33



Dye 34




5 Dyes 35 to 37



Dye No.	Z	An <sup>•</sup>
35	C <sub>6</sub> H <sub>5</sub>	NO <sub>3</sub> <sup>•</sup>
36	C <sub>2</sub> H <sub>5</sub>	NO <sub>3</sub> <sup>•</sup>
37	C <sub>6</sub> H <sub>5</sub>	BF <sub>4</sub> <sup>•</sup>


5

Dyes 38 to 42



Farb-stoff Nr.	Q <sup>1</sup>	Q <sup>2</sup>	Q <sup>3</sup>	Q <sup>4</sup>	An <sup>•</sup>
38	CH <sub>3</sub>	CH <sub>3</sub>	Cl	Cl	I <sup>•</sup>
39	CH <sub>3</sub>	CH <sub>3</sub>	H	H	I <sup>•</sup>
40	$\begin{array}{c} \text{O} \\ \parallel \\ \text{C}_2\text{H}_4\text{OCNHC}(\text{CH}_3)_3 \end{array}$	$\begin{array}{c} \text{O} \\ \parallel \\ \text{C}_2\text{H}_4\text{OCNHC}(\text{CH}_3)_3 \end{array}$	H	H	I <sup>•</sup>
41	$\begin{array}{c} \text{O} \\ \parallel \\ \text{C}_2\text{H}_4\text{CNHC}_6\text{H}_{13} \end{array}$	$\begin{array}{c} \text{O} \\ \parallel \\ \text{C}_2\text{H}_4\text{CNHC}_6\text{H}_{13} \end{array}$	H	H	ClO <sub>4</sub> <sup>•</sup>
42	C <sub>3</sub> H <sub>6</sub> SO <sub>3</sub> <sup>•</sup>	C <sub>3</sub> H <sub>6</sub> SO <sub>3</sub> H	H	H	Betain

Dye 43



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Use of a compound from the class of metal-free or metal-containing phthalocyanines, metal-free or metal-containing naphthalocyanines, nickel dithiolene complexes, aminium compounds of aromatic amines, methine dyes or azulenesquaric acid dyes which have their absorption maximum in the range from 600 to 1,200 nm and/or a fluorescence mixture in the range from 620 to 1,200 nm, as a marker for liquids.
2. A use as claimed in claim 1, wherein the compounds comes from the class of metal-free or metal-containing naphthalocyanines or nickel dithiolene complexes.
3. A use as claimed in claim 1, wherein the compound comes from the class of metal-free or metal-containing naphthalocyanines.
4. A method for detecting a marker from the class of metal-free or metal-containing naphthalocyanines, metal-free or metal-containing naphthalocyanines, nickel dithiolene complexes, aminium compounds of aromatic amines, methine dyes or azulenesquaric acid dyes which have their fluorescence maximum in the range from 620 to 1,200 nm by exciting its fluorescence and so producing fluorescent light in the NIR spectral region.
5. A method as claimed in claim 4, wherein the fluorescence is excited by means of a semiconductor laser or a semiconductor diode.
6. A method as claimed in claim 4, wherein the fluorescence light is detected by means of a semiconductor detector.
7. A method as claimed in claim 4, wherein a semiconductor laser or a semiconductor diode having a wavelength of maximum emission in the spectral range from  $\lambda_{\max}-100$  nm to  $\lambda_{\max}+20$  nm is used, where  $\lambda_{\max}$  is the wavelength of the absorption maximum of the marker.






8. A method as claimed in claim 4, wherein a semiconductor laser or a semiconductor diode having a wavelength of maximum emission of from 620 to 1,200 nm is used.


DATED this 27 day of June, 1996

B A S F AKTIENGESELLSCHAFT

WATERMARK PATENT & TRADEMARK ATTORNEYS  
290 BURWOOD ROAD  
HAWTHORN VICTORIA 3122  
AUSTRALIA

LCG/JGC/SH DOC 7 AU4567893.WPC





## A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl.<sup>5</sup> C10L 1/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl.<sup>5</sup> C10L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 4209302 (ORELUP, R.B.) 24 June 1980 (24.06.80) claim 1	1-3
A	US, A, 4009008 (ORELUP, R.B.) 22 February 1977 (22.02.77) abstract	1
A	US, A, 4735631 (ORELUP, R.B.) 5 April 1988 (05.04.88) abstract	1
A	EP, A2, 0155780 (IMPERIAL CHEMICAL INDUSTRIES PLC) 25 September 1985 (25.09.85) claim 1	1-3
A	EP, A2, 0358080 (BASF AKTIENGESELLSCHAFT) 14 March 1990 (14.03.90) claim 1	1-3
A	EP, A1, 0192215 (BASF AKTIENGESELLSCHAFT) 27 August 1986 (27.08.86) claim 1	1
A	US, A, 3484467 (SUSI, P.V. ET AL.) 16 December 1969 (16.12.69) claim 1	1
		./.

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

21 September 1993 (21.09.93)

Date of mailing of the international search report

29 October 1993 (29.10.93)

Name and mailing address of the ISA/

EUROPEAN PATENT OFFICE

Authorized officer

Facsimile No.

Telephone No.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 93/01830

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP, A1, 0464543 (BASF AKTIENGESELLSCHAFT) 8 January 1992 (08.01.92) claim 1 ---	1
A	EP, A2, 0310080 (BASF AKTIENGESELLSCHAFT) 5 April 1989 (05.04.89) claim 1  -----	1

## INTERNATIONAL SEARCH REPORT


Information on patent family members

26/08/93

International application No.

PCT/EP 93/01830

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4209302	24/06/80	BE-A- 883157	01/09/80
		CA-A- 1128513	27/07/82
		CH-A,B- 648587	29/03/85
		DE-A,C- 3017653	29/01/81
		FR-A,B- 2456322	05/12/80
		GB-A,B- 2050370	07/01/81
		NL-A- 8002708	12/11/80
US-A- 4009008	22/02/77	BE-A- 807508	20/05/74
		CA-A- 990716	08/06/76
		CH-A- 606306	31/10/78
		DE-A,C- 2355385	12/06/74
		FR-A,B- 2212390	26/07/74
		GB-A- 1394018	14/05/75
		JP-C- 1085216	25/02/82
		JP-C- 1146254	12/05/83
		JP-A- 49088925	26/08/74
		JP-A- 50142625	17/11/75
		JP-B- 56017390	22/04/81
		JP-B- 57036940	06/08/82
		LU-A- 68949	05/07/74
		NL-A- 7315382	11/06/74
		US-A- 3862120	21/01/75
		GB-A- 1477552	22/06/77
		US-A- 4049393	20/09/77
US-A- 4735631	05/04/88	CA-A- 1229595	24/11/87
		EP-A,B- 0147704	10/07/85
		EP-A,B- 0149125	24/07/85
		US-A- 4764474	16/08/88
EP-A2- 0155780	25/09/85	JP-B- 4075916	02/12/92
		JP-A- 60209583	22/10/85
		US-A- 4606859	19/08/86
EP-A2- 0358080	14/03/90	DE-A- 3830041	08/03/90
		JP-A- 2117964	02/05/90
		US-A- 5047312	10/09/91
EP-A1- 0192215	27/08/86	DE-A- 3505751	21/08/86
		DE-A- 3660154	01/06/88
		JP-B- 5011120	12/02/93
		JP-A- 61225192	06/10/86
		US-A- 4806664	21/02/89
US-A- 3484467	16/12/69	US-A- 3670025	13/06/72
EP-A1- 0464543	08/01/92	DE-A- 4021078	09/01/92
		JP-A- 4252268	08/09/92
EP-A2- 0310080	05/04/89	DE-A- 3733173	20/04/89
		JP-A- 1110562	27/04/89
		US-A- 5084592	28/01/92
		DE-A- 3816187	23/11/89

<b>A. KLASIFIZIERUNG DES ANMELDUNGSGEGENSTANDES</b>		
IPC5: C10L 1/10 Nach der Internationalen Patentklassifikation (IPK) oder nach der nationalen Klassifikation und der IPK		
<b>B. RECHERCHIERTE GEBIETE</b>		
Recherchierte Mindestprüfstoff (Klassifikationssystem und Klassifikationssymbole)		
IPC5: C10L		
Recherche, aber nicht zum Mindestprüfstoff gehorende Veröffentlichungen, soweit diese unter die recherchierten Gebiete fallen		
Während der internationalen Recherche konsultierte elektronische Datenbank (Name der Datenbank und evtl. verwendete Suchbegriffe)		
WPI, CLAIMS/US PATENT ABS, JAPIO, CA SEARCH		
<b>C. ALS WESENTLICH ANGESEHENE UNTERLAGEN</b>		
Kategorie*	Bezeichnung der Veröffentlichung, soweit erforderlich unter Angabe der in Betracht kommenden Teile	Betr. Anspruch Nr.
X	US, A, 4209302 (ORELUP, R.B.), 24 Juni 1980 (24.06.80), Anspruch 1 --	1-3
A	US, A, 4009008 (ORELUP, R.B.), 22 Februar 1977 (22.02.77), Zusammenfassung --	1
A	US, A, 4735631 (ORELUP, R.B.), 5 April 1988 (05.04.88), Zusammenfassung --	1
A	EP, A2, 0155780 (IMPERIAL CHEMICAL INDUSTRIES PLC), 25 September 1985 (25.09.85), Anspruch 1 --	1-3
<input checked="" type="checkbox"/> Weitere Veröffentlichungen sind der Fortsetzung von Feld C zu entnehmen. <input checked="" type="checkbox"/> Siehe Anhang Patentfamilie.		
* Besondere Kategorien von angegebenen Veröffentlichungen: "A" Veröffentlichung, die den allgemeinen Stand der Technik definiert, aber nicht als besonders bedeutsam anzusehen ist "E" älteres Dokument, das jedoch erst am oder nach dem internationalen Anmeldedatum veröffentlicht worden ist "L" Veröffentlichung, die geeignet ist, einen Prioritätsanspruch zweifelhaft erscheinen zu lassen, durch die das Veröffentlichungsdatum einer späteren im Recherchenbericht genannten Veröffentlichung belegt werden soll oder die aus einem anderen besonderen Grund angegeben ist (wie ausgeführt) "O" Veröffentlichung, die sich auf eine mündliche Offenbarung, eine Benutzung, eine Ausstellung oder andere Maßnahmen bezieht "P" Veröffentlichung, die vor dem internationalen Anmeldedatum, aber nach dem beanspruchten Prioritätsdatum veröffentlicht worden ist "T" Spätere Veröffentlichung, die nach dem internationalen Anmeldedatum oder dem Prioritätsdatum veröffentlicht worden ist und mit der Anmeldung nicht zusammenfällt, sondern nur zum Verständnis der der Erfindung zugrundeliegenden Prinzipien oder der ihr zugrundeliegenden Theorie angegeben ist "X" Veröffentlichung von besonderer Bedeutung: die beanspruchte Erfindung kann allein aufgrund dieser Veröffentlichung nicht als neu oder auf erfindungsmäßiger Tätigkeit beruhend betrachtet werden "Y" Veröffentlichung von besonderer Bedeutung: die beanspruchte Erfindung kann nicht als auf erfindungsmäßiger Tätigkeit beruhend betrachtet werden, wenn die Veröffentlichung mit einer oder mehreren Veröffentlichungen dieser Kategorie in Verbindung gebracht wird und diese Verbindung für einen Fachmann absehbar ist "Z" Veröffentlichung, die Mitglied derselben Patentfamilie ist		
Datum des Abschlusses der internationalen Recherche		Absenddatum des internationalen Recherchenberichts
21 September 1993		29.10.93
Name und Postanschrift der Internationalen Recherchenbehörde		Bevollmächtigter Beauftragter
 Europäisches Patentamt, P.B. 5818 Patentlaan 2 NL-2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016		INGA-KARIN PETERSSON

## C (Fortsetzung). ALS WESENTLICH ANGESEHENE UNTERLAGEN

Kategorie*	Bezeichnung der Veröffentlichung, soweit erforderlich unter Angabe der in Betracht kommenden Teile	Bez. Anspruch Nr.
A	EP, A2, 0358080 (BASF AKTIENGESELLSCHAFT), 14 März 1990 (14.03.90), Anspruch 1 --	1-3
A	EP, A1, 0192215 (BASF AKTIENGESELLSCHAFT), 27 August 1986 (27.08.86), Anspruch 1 --	1
A	US, A, 3484467 (SUSI, P.V. ET AL.), 16 Dezember 1969 (16.12.69), Anspruch 1 --	1
A	EP, A1, 0464543 (BASF AKTIENGESELLSCHAFT), 8 Januar 1992 (08.01.92), Anspruch 1 --	1
A	EP, A2, 0310080 (BASF AKTIENGESELLSCHAFT), 5 April 1989 (05.04.89), Anspruch 1 -- -----	1

## INTERNATIONALER RECHERCHENBERICHT

Angaben zu Veröffentlichungen, die zur selben Patentfamilie gehören  
26/08/93

Internationales Aktenzeichen

PCT/EP 93/01830

Im Recherchenbericht angeführtes Patentdokument	Datum der Veröffentlichung	Mitglied(er) der Patentfamilie	Datum der Veröffentlichung
US-A- 4209302	24/06/80	BE-A- 883157	01/09/80
		CA-A- 1128513	27/07/82
		CH-A,B- 648587	29/03/85
		DE-A,C- 3017653	29/01/81
		FR-A,B- 2456322	05/12/80
		GB-A,B- 2050370	07/01/81
		NL-A- 8002708	12/11/80
US-A- 4009008	22/02/77	BE-A- 807508	20/05/74
		CA-A- 990716	08/06/76
		CH-A- 606306	31/10/78
		DE-A,C- 2355385	12/06/74
		FR-A,B- 2212390	26/07/74
		GB-A- 1394018	14/05/75
		JP-C- 1085216	25/02/82
		JP-C- 1146254	12/05/83
		JP-A- 49088925	26/08/74
		JP-A- 50142625	17/11/75
		JP-B- 56017390	22/04/81
		JP-B- 57036940	06/08/82
		LU-A- 68949	05/07/74
		NL-A- 7315382	11/06/74
		US-A- 3862120	21/01/75
		GB-A- 1477552	22/06/77
		US-A- 4049393	20/09/77
US-A- 4735631	05/04/88	CA-A- 1229595	24/11/87
		EP-A,B- 0147704	10/07/85
		EP-A,B- 0149125	24/07/85
		US-A- 4764474	16/08/88
EP-A2- 0155780	25/09/85	JP-B- 4075916	02/12/92
		JP-A- 60209583	22/10/85
		US-A- 4606859	19/08/86
EP-A2- 0358080	14/03/90	DE-A- 3830041	08/03/90
		JP-A- 2117964	02/05/90
		US-A- 5047312	10/09/91
EP-A1- 0192215	27/08/86	DE-A- 3505751	21/08/86
		DE-A- 3660154	01/06/88
		JP-B- 5011120	12/02/93
		JP-A- 61225192	06/10/86
		US-A- 4806664	21/02/89
US-A- 3484467	16/12/69	US-A- 3670025	13/06/72
EP-A1- 0464543	08/01/92	DE-A- 4021078	09/01/92
		JP-A- 4252268	08/09/92
EP-A2- 0310080	05/04/89	DE-A- 3733173	20/04/89
		JP-A- 1110562	27/04/89
		US-A- 5084592	28/01/92
		DE-A- 3816187	23/11/89